



Trends in the Prevalence of Tobacco Use

National YRBS: 1991–2009

The national Youth Risk Behavior Survey (YRBS) monitors priority health risk behaviors that contribute to the leading causes of death, disability, and social problems among youth and adults in the United States. The national YRBS is conducted every two years during the spring semester and provides data representative of 9th through 12th grade students in public and private schools throughout the United States.

1991	1993	1995	1997	1999	2001	2003	2005	2007	2009	Changes from 1991 2009 ¹	Change from 2007 2009 ²
Ever tried cigarette smoking (even one or two puffs)											
70.1 (67.8–72.3) ³	69.5 (68.1–70.8)	71.3 (69.5–73.0)	70.2 (68.2–72.1)	70.4 (67.3–73.3)	63.9 (61.6–66.0)	58.4 (55.1–61.6)	54.3 (51.2–57.3)	50.3 (47.2–53.5)	46.3 (43.7–48.9)	No change, 1991–1999 Decreased, 1999–2009	Decreased
Smoked cigarettes on at least 1 day (during the 30 days before the survey)											
27.5 (24.8–30.3)	30.5 (28.6–32.4)	34.8 (32.5–37.2)	36.4 (34.1–38.7)	34.8 (32.3–37.4)	28.5 (26.4–30.6)	21.9 (19.8–24.2)	23.0 (20.7–25.5)	20.0 (17.6–22.6)	19.5 (17.9–21.2)	Increased, 1991–1997 Decreased, 1997–2009	No change
Smoked cigarettes on 20 or more days (during the 30 days before the survey)											
12.7 (10.6–15.3)	13.8 (12.1–15.5)	16.1 (13.6–19.1)	16.7 (14.8–18.7)	16.8 (14.3–19.6)	13.8 (12.3–15.5)	9.7 (8.3–11.3)	9.4 (7.9–11.0)	8.1 (6.7–9.8)	7.3 (6.4–8.3)	Increased, 1991–1999 Decreased, 1999–2009	No change
Used chewing tobacco, snuff, or dip on at least 1 day (during the 30 days before the survey)											
NA ⁴	NA	11.4 (9.8–13.2)	9.3 (7.3–11.8)	7.8 (5.8–10.4)	8.2 (6.8–9.9)	6.7 (5.3–8.5)	8.0 (6.6–9.6)	7.9 (6.3–9.8)	8.9 (7.3–10.8)	Decreased, 1995–2003 No change, 2003–2009	No change
Smoked cigars, cigarillos, or little cigars on at least 1 day (during the 30 days before the survey)											
NA	NA	NA	22.0 (19.9–24.2)	17.7 (16.1–19.5)	15.2 (14.0–16.5)	14.8 (13.2–16.7)	14.0 (12.6–15.6)	13.6 (12.1–15.2)	14.0 (12.8–15.4)	Decreased, 1997–2005 No change, 2005–2009	No change
Smoked cigarettes; smoked cigars, cigarillos, or little cigars; or used chewing tobacco, snuff, or dip on at least 1 day (during the 30 days before the survey)											
NA	NA	NA	43.4 (41.0–45.8)	40.2 (37.4–43.0)	33.9 (31.8–36.1)	27.5 (25.1–30.0)	28.4 (25.7–31.3)	25.7 (22.8–28.7)	26.0 (23.8–28.3)	Decreased, 1997–2003 No change, 2003–2009	No change

¹ Based on trend analyses using a logistic regression model controlling for sex, race/ethnicity, and grade.

² Based on t-test analyses, $p < 0.05$.

³ 95% confidence interval.

⁴ Not available.

Where can I get more information? Visit www.cdc.gov/healthyyouth or call 800–CDC–INFO (800–232–4636).



U.S. Department of Health and Human Services

Centers for Disease Control and Prevention

National Center for Chronic Disease Prevention and Health Promotion

Division of Adolescent and School Health

